arXiv:1205.2591v2 [cond-mat.stat-mech] 21 Jul 2012

Scalar curvature of systems with fractal distribution functions

Marcelo R. Ubriaco*

Laboratory of Theoretical Physics
Department of Physics
University of Puerto Rico
Río Piedras Campus
San Juan
PR 00931, USA

Abstract

Starting with the relative entropy for two close statistical states we define the metric and calculate the scalar curvature R for systems with classical, boson and fermion fractal distribution functions with moment order parameter q. In particular, we find that for $q \neq 1$ the scalar curvature is closer to zero implying that the fractal bosonic and fermionic systems are more stable than the standard ones.

Keywords: Scalar Curvature, fractal systems, stability
PACS:05.20.-y, 02.40.-k, 02.50.-r, 05.45.Df

1 Introduction

From the theory of fractals [1] we learned that given a statistical weight $\Omega(q,\delta)$ of a system with order parameter q and resolution δ, the fractal dimension is defined as the exponent $d = D_q$ which will make the product $lim_{\delta\to 0}\Omega(q,\delta)\delta^d$ finite. With use of the definition of the Boltzmann entropy $S(q,\delta) = \ln\Omega(q,\delta)$, the relation between the entropy and the fractal dimension D_q is given by

$$D_q = -\lim_{\delta\to 0}\frac{S(q,\delta)}{\ln\delta}. \tag{1}$$

Based on these definitions and with use of the Boltzmann's H theorem, the generalized entropy and distribution functions for classical and quantum gases

---

*Electronic address:ubriaco@ltp.uprrp.edu

were calculated in Ref.[2]. The average number of particles with energy ϵ was shown to be given by

$$< n(\epsilon) >= \frac{1}{[1+\beta(q-1)(\epsilon-\mu)]^{1/(q-1)}+a}, \tag{2}$$

where $a = 0$ for the classical case, and the values $a = -1$ and $a = 1$ correspond to Bose-Einstein and Fermi-Dirac cases, respectively. For $q = 1$, Equation (2) becomes the standard textbook result for classical and quantum ideal gases. The distribution functions in Equation (2) were also obtained in Ref.[3] by considering a dilute gas approximation to the partition function of a non-extensive statistical mechanics originally proposed in Ref.[4]. It is our purpose to study some of the geometric properties of systems with average particle number according to Equation (2). The idea of using geometry in thermodynamics is not new [5]-[9], and several authors developed formalisms to measure the distance between equilibrium states through the definition of a metric and the calculation of the corresponding scalar curvature as a measure of the interactions [10]-[20]. Some of the applications include classical and quantum gases [11][16][21][22], magnetic systems [23]-[26], non-extensive statistical thermodynamics [27][28][29], anyon gas [30][31], fractional statistics [32] and deformed boson and fermion systems [33]. Some of the basic results of these approaches include the relationships between the metric with the correlations of the stochastic variables, and the scalar curvature R with the stability of the system, and the facts that the scalar curvature R vanishes for the classical ideal gas, $R > 0 (R < 0)$ for a boson (fermion) ideal gas, and it is singular at a critical point. Here, we wish to study systems with an average particle number given in Equation (2). In Section 2 we briefly describe the formalism of systems with fractal distribution functions as reported in Ref. [34]. In Section 3 we obtain the metric from the second order term in the expansion of the relative entropy between two close statistical states, and in Section 4 we use the metric to compute the scalar curvature for the classical, Bose-Einstein and Fermi-Dirac cases for $q \neq 1$. In Section 5 we summarize our results.

2 Fractal models

2.1 Classical case

We will use the short notation

$$\rho_l = [1+(q-1)\beta(\epsilon_l-\mu)]^{1/(q-1)}. \tag{3}$$

The probability density is defined

$$\rho = \frac{1}{Z_{MB}} \prod_{l=0} \frac{1}{n_l!} \rho_l^{-n_l}, \tag{4}$$

where the partition function

$$\begin{aligned} Z_{MB} &= \prod_{l=0} \sum_{n_l=0} \frac{1}{n_l!} \rho_l^{-n_l} \\ &= \prod_{l=0} e^{\rho_l^{-1}} \end{aligned} \tag{5}$$

From the definition of the average number of particles with energy ϵ_l

$$< n_l >= \frac{\sum_{n_l=0} \frac{1}{n_l!} n_l \rho_l^{-n_l}}{\sum_{n_l=0} \frac{1}{n_l!} \rho_l^{-n_l}}, \tag{6}$$

we find after summing the series that

$$< n_l >= \rho_l^{-1}. \tag{7}$$

as required. In the thermodynamic limit we write for the average total number of particles $< N >$

$$< N >= \frac{4\pi V}{h^3} \left(\frac{2m}{\beta(q-1)} \right)^{3/2} \int_0^\infty \frac{x^2 dx}{[1 + x^2 - (q-1)\beta\mu]^{1/(q-1)}}, \tag{8}$$

leading to the expression

$$< N >= \frac{-2\pi V}{h^3} \left(\frac{2m}{\beta(q-1)} \right)^{3/2} \frac{1}{[1 - (q-1)\beta\mu]^{1/(q-1)-(3/2)}} S, \tag{9}$$

where S is a series independent of $\beta\mu$, given by

$$S = C_0 + \sum_{i=1}^{\infty} \frac{(-1)^m}{m!} \left(\frac{1}{2} \right) \ldots \left(\frac{3}{2} - m \right) C_m, \tag{10}$$

with $C_m = \frac{1}{-1/(q-1)-m+(3/2)}$. Solving Equation (8) we find that the fugacity $z = e^{\beta\mu}$ has a temperature dependence given by

$$\ln z = \frac{1}{q-1} \left\{ 1 - \left[\frac{-2\pi V}{h^3} \left(\frac{2m}{(q-1)\beta} \right)^{3/2} S \right]^{1/\omega} \right\}, \tag{11}$$

where $\omega = \frac{5-3q}{2(q-1)}$. From Equation (11) we see that the fugacity is restricted to the interval $0 < z < e^{1/(q-1)}$, which serves as a cut-off that avoids a negative average occupation number. The correct definition of the average energy is given by

$$< \epsilon >= \frac{4\pi V}{h^3} \int_0^\infty \frac{p^2}{2m} < n(p) >^q p^2 dp, \tag{12}$$

leading, with Equation (8), to the required classical result $< \epsilon >= \frac{3}{2} < N > kT$ [34]. We should remark, that Equations (5) and (12) are related by the standard definition

$$< \epsilon >= -\frac{\partial \ln Z_{MB}}{\partial \beta}. \tag{13}$$

2.2 Boson case

Similarly to the classical case, the definition

$$< N >= \sum_{j=0} \frac{\sum_{n_j=0} n_j \rho_j^{-n_j}}{\sum_{n_j=0} \rho_j^{-n_j}}, \tag{14}$$

leads to the average occupation number

$$< n_j >= \frac{1}{\rho_l - 1} \tag{15}$$

with the probability density and the partition function [1]

$$\begin{aligned} \rho &= \frac{1}{Z_{BE}} \prod_{j=0} \rho_j^{-n_j}, & (16) \\ Z_{BE} &= \prod_{j=0} \sum_{n_j=0} \rho_j^{-n_j} \\ &= \prod_{j=0} \frac{1}{1-\rho_j^{-1}}. & (17) \end{aligned}$$

As in the standard, $q = 1$, Bose-Einstein case the chemical potential is negative.

2.3 Fermion case

For the Fermi-Dirac case the average occupation number

$$< n_j >= \frac{1}{\rho_l + 1} \tag{18}$$

is obtained by defining

$$\begin{aligned} \rho &= \frac{1}{Z_{FD}} \prod_{j=0} \rho_j^{-n_j}, & (19) \\ Z_{FD} &= \prod_{j=0} \sum_{n_j=0}^{1} \rho_j^{-n_j} \\ &= \prod_{j=0} (1+\rho_j^{-1}), & (20) \end{aligned}$$

and the requirement that the average occupation number $< n_j > \in [0, 1]$ leads to restrict the fugacity to the interval $0 < z < e^{1/(q-1)}$. For Bose-Einstein and Fermi-Dirac cases we obtain

$$-\frac{\partial \ln Z}{\partial \beta} =< \epsilon > + \sum_{l=0} \sum_{k=1}^{\infty} \left(\begin{array}{c} 1-q \\ k \end{array} \right) < n_l >^{q+k} \epsilon_l (-a)^k. \tag{21}$$

---

[1] We should remark that our partition function differs from that reported in Ref. [3].

3 The metric

For the three cases discussed in the previous section we cannot adopt any of the standard definitions for the metric like for example [21]

$$g_{\alpha\gamma} = \frac{\partial^2 \ln Z}{\partial\beta^\alpha \partial\beta^\gamma}, \quad \beta^1 = \beta \ ; \ \beta^2 = -\beta\mu \tag{22}$$

which is valid for exponential distributions.
The relative entropy, $H(p||P) = \sum_i p_i \left(-\ln(\frac{P_i}{p_i}) \right)^\mu$, is a very useful concept. For example, for two close distribution functions $p(x)$ and $p(x+\Delta)$ we can obtain a Fisher's information measure

$$I_\mu = \int dx \left(\frac{dp/dx}{p(x)} \right)^\mu \frac{dp}{dx}, \tag{23}$$

as part of the second order term in Δ for the entropic form, with $\mu = 1$, $S = -\sum_i p_i \ln p_i$ [35], and $S = \sum_i p_i(-\ln p_i)^\mu$ [36] where μ is a fractional parameter. By defining $\phi(x) = p^{\frac{1}{1+\mu}}$ and considering the Fisher information as a lagrangian density leads to linear and nonlinear differential equations for $\mu = 1$ and $\mu \neq 1$, respectively.

Here, based on work in Ref.[10] we expand the relative entropy for $\mu = 1$ between two close densities $\rho(\beta)$ and $\rho(\beta + d\beta)$ up to second order in $d\beta^\alpha$. Therefore, the information distance $I(\rho(\beta), \rho(\beta + d\beta))$ between the two close states is written

$$I(\rho(\beta^\alpha), \rho(\beta^\alpha + d\beta^\alpha)) = Tr\rho \left(\ln \rho(\beta^\alpha) - \ln \rho(\beta^\alpha + d\beta^\alpha) \right), \tag{24}$$

such that expanding the second order term gives for the metric

$$g_{\alpha\gamma} = \frac{\partial^2 \ln Z}{\partial\beta^\alpha \partial\beta^\gamma} + Tr\rho \sum_{l=0} < n_l > \frac{\partial^2 \ln \rho_l}{\partial\beta^\alpha \partial\beta^\gamma}. \tag{25}$$

A simple inspection shows that in the $q \to 1$ limit Equation (25) reduces to Equation (22). Equation (25) can be simplified leading to the three corresponding metrics:

$$\begin{aligned} g^{MB}_{\alpha\gamma} &= \sum_{l=0} \frac{1}{\rho_l^3} \frac{\partial\rho_l}{\partial\beta^\alpha} \frac{\partial\rho_l}{\partial\beta^\gamma}, && (26) \\ g^{BE}_{\alpha\gamma} &= \sum_{l=0} \frac{1}{\rho_l(\rho_l - 1)^2} \frac{\partial\rho_l}{\partial\beta^\alpha} \frac{\partial\rho_l}{\partial\beta^\gamma}, && (27) \\ g^{FD}_{\alpha\gamma} &= \sum_{l=0} \frac{1}{\rho_l(\rho_l + 1)^2} \frac{\partial\rho_l}{\partial\beta^\alpha} \frac{\partial\rho_l}{\partial\beta^\gamma}, && (28) \\ & && (29) \end{aligned}$$

which can be summarized in the general formula

$$g_{\alpha\gamma} = \sum_{l=0} \frac{< n_l >^2}{\rho_l} \frac{\partial\rho_l}{\partial\beta^\alpha} \frac{\partial\rho_l}{\partial\beta^\gamma}. \tag{30}$$

Writing in general $\rho_l = [1 + (q-1)\sum_\alpha \beta^\alpha F_l^\alpha]^{1/(q-1)}$ we find

$$\begin{aligned} \frac{\partial}{\partial\beta^\alpha}\frac{\partial}{\partial\beta^\lambda}\ln Z &= \sum_{l=0} \rho_l^{2-2q} F_l^\alpha F_l^\lambda \left(q < n_l > -a < n_l >^2\right) && (31) \\ g_{\alpha\lambda} &= \sum_{l=0} \rho_l^{2-2q} F_l^\alpha F_l^\lambda \left(< n_l > -a < n_l >^2\right) && (32) \end{aligned}$$

4 Scalar curvature

4.1 Classical case

It has been shown that the scalar curvature [11][16] vanishes for the standard case, but it is tempting to speculate whether that is also the case for $q \neq 1$. In the thermodynamic limit,with $x = \beta\epsilon$, we write for example

$$g_{11} = \frac{2}{\sqrt{\pi}} V\beta^{-2}\lambda^{-3} \int_0^\infty \frac{x^{5/2}dx}{[1 + (q-1)(x+\gamma)]^{\frac{2q-1}{q-1}}} \qquad (33)$$

where hereafter $\gamma = -\beta\mu$. This integral converges for $\frac{5-3q}{2(q-1)} > 0$, restricting the values of q to the interval $q \in [1, 5/3)$. With use of the integral representation of the Γ-function

$$\Gamma(y) = w^y \int_0^\infty t^{y-1} e^{-wt} dt, \ \ y > 0 \ ; \ w > 0, \qquad (34)$$

we obtain for the components of the metric tensor

$$\begin{aligned} g_{11} &= V\beta^{-2}\lambda^{-3} h_{5/2}, \\ g_{12} &= V\beta^{-1}\lambda^{-3} h_{3/2}, \\ g_{22} &= V\lambda^{-3} h_{1/2}, \end{aligned} \qquad (35)$$

where the function

$$h_\lambda = \frac{2}{\sqrt{\pi}(q-1)^{(\lambda+1)}} \frac{\Gamma(\lambda+1)\Gamma(\frac{q}{q-1} - \lambda)}{\Gamma(\frac{2q-1}{q-1})} \frac{1}{[1 + (q-1)\gamma]^{\frac{q}{q-1} - \lambda}}, \qquad (36)$$

satisfies

$$\frac{\partial h_\lambda}{\partial\gamma} = -\lambda h_{\lambda-1}. \qquad (37)$$

As is well known [17], the scalar curvature is given by

$$R = \frac{2}{detg} R_{1212}, \qquad (38)$$

where $detg = g_{11}g_{22} - g_{12}g_{12}$ and the non-vanishing part of the curvature tensor $R_{\alpha\beta\gamma\lambda}$ is given in terms of the Christoffel symbols



Figure 1: The scalar curvature R, in units of $\lambda^3 V^{-1}$, as a function of the fugacity z for bosons at constant β for the cases of $q = 1$ (solid line), $q = 1.1$ (dashed line) and $q = 1.2$ (dotted line).

$$R_{\alpha\beta\gamma\lambda} = g^{\eta\theta} \left(\Gamma_{\eta\alpha\lambda}\Gamma_{\theta\beta\gamma} - \Gamma_{\eta\alpha\gamma}\Gamma_{\theta\beta\lambda} \right). \tag{39}$$

A simple calculation leads to the result

$$R = \frac{V^{-1}\lambda^3}{4(detg)^2} \left(5h_{1/2}h_{3/2}^2 - 6h_{1/2}^2 h_{5/2} + h_{3/2}h_{-1/2}h_{5/2} \right), \tag{40}$$

such that after replacement of the definition of the function h_λ we get that the scalar curvature for the classical fractal case is identically equal to zero. Therefore, in this case the parameter q does not play any role as far as correlations are concerned.

4.2 Boson and fermion cases

Here, in order to evaluate the corresponding scalar curvatures we need to replace the summations in Equations (27) and (28) by integrals

$$G_\lambda^\pm = \frac{2}{\sqrt{\pi}} \int_0^\infty \frac{x^\lambda \Omega^{\frac{3-2q}{q-1}}}{(\Omega^{\frac{1}{q-1}} \pm 1)^2} dx, \tag{41}$$

where the + sign is for fermions and the − sign for bosons, and the function $\Omega = 1 + (q-1)(x+\gamma)$. In particular, the metric component g_{11} is written

$$g_{11} = V\lambda^{-3}\beta^{-2}G_{5/2}^\pm, \tag{42}$$

and its integral converges for $1 \leq q < 5/3$.

The functions $G_\lambda^\pm$ also satisfy

$$\frac{\partial G_\lambda^\pm}{\partial \gamma} = -\lambda G_{\lambda-1}^\pm, \tag{43}$$



Figure 2: The scalar curvature R, in units of $\lambda^3 V^{-1}$, as a function of the fugacity z for fermions at constant β for the cases of $q = 1$ (solid line), $q = 1.1$ (dashed line) and $q = 1.2$ (dotted line).

and thus the corresponding equations for R are equivalent to Equation (40) with the replacement of the function h_λ by the functions $G^{\pm}_{\lambda}$. Figures 1 and 2 show the results of a numerical calculation of the scalar curvature R as a function of the fugacity z for the parameter values $q = 1, 1.1, 1.2$ for boson and fermions respectively.

5 Conclusions

In this paper, starting from the relative entropy for two close statistical states we defined the metric for systems with fractal distribution functions with order parameter q. We calculated the scalar curvature R and found that it vanishes for the classical ideal gas, as in the standard case. Numerical calculations for the boson and fermion systems show that the corresponding values of R as a function of the fugacity z are closer to zero than those in the $q = 1$ case, implying that the departure from the value $q = 1$ makes the systems more stable. Therefore, for $q \neq 1$ bosons will be less attractive and fermions less repulsive that their standard counterparts. Our results are in agreement with those obtained in a previous work [37] wherein we showed that long-range correlations for the fractal Bose case decrease when the parameter q departs from the standard value $q = 1$. On the other hand, if one wishes to consider the order parameter q as a non-extensive parameter it has to be within the context of considering these fractal systems as a dilute approximation to non-extensive statistical mechanics, which consists in replacing the Tsallis partition function by a factorized one. This type of approximation has been shown [38] to be good outside a temperature interval that shifts to higher values of T when the number of energy levels increases. Our results also show that the sign of R remains unchanged as a function of z implying that these systems do not exhibit anyonic behavior, a fact that looks impossible to check by performing an expansion for $z \approx 0$ to obtain the second virial coefficient because the partition function is a function of $\ln z$. In addition, our results contrast with the cases of systems with quantum group symmetry where the parameter q interpolates between bosons and fermions in two and three dimensions [39].

Acknowledgements

I am grateful to the anonymous reviewers for their comments and constructive criticism to improve the original manuscript.

References

[1] J. Feder, Fractals (Plenum, New York, 1988).

[2] F. Büyükkiliç and D. Demirhan, Phys. Lett. A 181 (1993) 24.

[3] F. Büyükkiliç, D. Demirhan and A. Güleç, Phys. Lett. A 197 (1995) 209.

[4] C. Tsallis, J. Stat. Phys. 52 (1988) 479.

[5] L. Tisza, Generalized Themodynamics (MIT, Cambridge, 1966).

[6] R. B. Griffiths and J. C. Wheeler, Phys. Rev. A 2 (1970) 1047.

[7] F. Weinhold, J. Chem. Phys. 63 (1975) 2479.

[8] S.-I. Amari, Differential-Geometrical Methods in Statistics (Springer-Verlag, Berlin, 1985).

[9] S.-I. Amari and H. Nagaoka, Methods of Information Geometry (AMS, Rhode Island, 2000).

[10] R. S. Ingarden, H. Janyszek, A. Kossakowski and T. Kawaguchi, Tensor N.S. 37 (1982) 105.

[11] G. Ruppeiner, Phys. Rev. A 20 (1979) 1608.

[12] W. K. Wootters, Phys. Rev. D 23 (1981) 357.

[13] R. Gilmore, Phys. Rev. A 30 (1984) 1994.

[14] G. Ruppeiner, Phys. Rev. A 32 (1985) 3141.

[15] R. Gilmore, Phys. Rev. A 32 (1985) 3144.

[16] J. Nulton, P. Salamon, Phys. Rev. A 31 (1985) 2520.

[17] S. Weinberg, Gravitation and Cosmology (Wiley and Sons, Inc., New York, 1972).

[18] H. Janyszek, Rep. Math. Phys. 24 (1986) 1; Rep. Math. Phys. 24 (1986)11.

[19] H. Janiszek and R Mrugala, Rep. Math. Phys. 27 (1989) 145.

[20] G. Ruppeiner, Am. J. Phys. 78 (2010) 1170, and references therein.

[21] H. Janiszek and R Mrugala, J. Phys. A: Math. Theor. 23 (1990) 467.

[22] D. Brody and D. Hook , J. Phys. A: Math. Theor.42 (2009) 023001.

[23] H. Janiszek and R Mrugala, Phys. Rev. A 39 (1989) 6515.

[24] H. Janyszek, J. Phys. A:Math. 23 (1990) 477.

[25] D. Brody and N. Rivier, Phys. Rev. E 51 (1995) 1006.

[26] W. Janke, D. A. Johnston and R. Kenna, Physica A 336 (2004) 181.

[27] R. Trasarti-Battistoni, cond-mat/0203536.

[28] M. Portesi, A. Plastino and F. Pennini, Physica A 365 (2006) 173.

[29] A. Ohara, Phys. Lett. A 370 (2007) 184.

[30] B. Mirza and H. Mohammadzadeh, Phys. Rev. E 79 (2008) 021127.

[31] B. Mirza and H. Mohammadzadeh, Phys. Rev. E 80 (2009) 011132.

[32] B. Mirza and H. Mohammadzadeh, Phys. Rev. E 82 (2010) 031137.

[33] B. Mirza and H. Mohammadzadeh ,J. Phys. A: Math. Theor. 44 (2011) 475003

[34] M. R. Ubriaco, Phys. Rev. E 60 (1999) 165.

[35] G.V Vstovsky, Phys. Rev. E 51 (1995) 975.

[36] M. R. Ubriaco, Phys. Lett. A 373 (2009) 4017.

[37] M. R. Ubriaco, Phys. Rev. E 62 (2000) 328.

[38] Q.-A. Wang and A. Le Méhauté, Phys. Lett. A 235 (1997) 222.

[39] M. R. Ubriaco, Phys. Rev. E 55 (1997) 291.